

DIDACTICS OF MATHEMATICS

12(16)



The Publishing House
of Wrocław University of Economics
Wrocław 2015

Reviewers cooperating with the journal

*Maria Balcerowicz-Szkutnik, Giovanna Carcano, Igor Dubina,
Ewa Dziwok, Salvatore Federico, Marian Matłoka,
Włodzimierz Odyniec, Anatol Pilawski,
Tadeusz Stanisz, Achille Vernizzi, Henryk Zawadzki*

Copy-editing

Elżbieta Macauley, Tim Macauley

Proof reading

Marcin Orszulak

Typesetting

Elżbieta Szlachcic

Cover design

Robert Mazurczyk

Front cover painting: W. Tank, Sower
(private collection)

This publication is available at: www.journal.ue.wroc.pl and www.ibuk.pl
BazEkon, http://kangur.uek.krakow.pl/bazy_ae/bazekon/nowy/advanced.php Dolnośląska
Biblioteka Cyfrowa, <http://www.dbc.wroc.pl/>
The Central European Journal of Social Sciences and Humanities, <http://cejsh.icm.edu.pl/>
Ebsco, <https://www.ebscohost.com/>

Information on submitting and reviewing papers is available on
the Publishing House's websites
www.dm.ue.wroc.pl; www.wydawnictwo.ue.wroc.pl

The publication is distributed under the Creative Commons Attribution 3.0
Attribution-NonCommercial-NoDerivs CC BY-NC-ND



© Copyright by Wrocław University of Economics
Wrocław 2015

ISSN 1733-7941
e-ISSN 2450-1123

The original version: printed
Publication may be ordered in Publishing House tel./
fax 71 36-80-602; e-mail: econbook@ue.wroc.pl
www.ksiegarnia.ue.wroc.pl
Printing: TOTEM

TABLE OF CONTENTS

Marek Biernacki	
Elements of differential equations in the mathematics course for students of economics	5
Marek Biernacki, Andrzej Misztal	
Is an average Polish student proficient in solving difficult and new problems?.....	11
Piotr Dniestrzański	
The Gini coefficient as a measure of disproportionality.....	25
Renata Dudzińska-Baryła, Donata Kopańska-Bródka, Ewa Michalska	
Software tools in didactics of mathematics	35
Ewa Dziwok	
The implementation of a double degree in Poland and its consequences for teaching quantitative courses	47
Wiktor Ejsmont	
Remarks on Wigner's semicircle law	55
Barbara Fura, Marek Fura	
Optimization of consumer preferences – an example.....	61
Donata Kopańska-Bródka, Renata Dudzińska-Baryła, Ewa Michalska	
An evaluation of the selected mathematical competence of the first-year students of economic studies	69
Arkadiusz Maciuk, Antoni Smoluk	
Two proofs of Stokes' theorem in new clothes	85
Paweł Prysak	
Mathematical preparation of first-year students of applied informatics for studies at the university of economics	93
Leszek Rudak	
“At 100 percent” assessment	111
Leszek Rudak, Mariusz Szalański	
Small Project Based Learning in a course of financial mathematics. A case study.....	117
Anna Szymańska, Elżbieta Zalewska	
E-learning as a tool to improve the quality of education in quantitative subjects	125
*	
Antoni Smoluk, Elżbieta Szlachcic	
Doktor inżynier Jerzy Sacała (1962-2015)	135

REMARKS ON WIGNER'S SEMICIRCLE LAW

Wiktor Ejsmont

Abstract. Ejsmont [2014] has shown that families of free Meixner distributions can be characterized by the conditional moments of polynomial functions of degree 3. In this paper, we will give other characterizations of the free normal distribution which are formulated in a similar spirit.

Keywords: Wigner's semicircle law, conditional expectation, free cumulants, Laha-Lukacs theorem, noncommutative regression.

JEL Classification: C10.

DOI: 10.15611/dm.2015.12.06.

1. Free probability, free cumulants, conditional expectation

We assume that our probability space is a von Neumann algebra A with a normal faithful tracial state $\tau: A \rightarrow C$ i.e., $\tau(\cdot)$ is linear, continuous in weak* topology, $\tau(XY) = \tau(YX)$, $\tau(1) = 1$, $\tau(XX^*) \geq 0$ and $\tau(XX^*) = 0$ implies $X = 0$ for all $X, Y \in A$. A (noncommutative) random variable V is a self-adjoint (i.e. $X = X^*$) element of A . The *-distribution μ of a self-adjoint element $X \in A$ is a probabilistic measure on R such that for all $n \geq 0$

$$\tau(X^n) = \int_R x^n d\mu(x).$$

Let $C\langle X_1, \dots, X_n \rangle$ denote the non-commutative ring of polynomials in variables X_1, \dots, X_n . The free cumulants are the k -linear maps $R_k: C\langle X_1, \dots, X_k \rangle \rightarrow C$ defined by the recursive formula (connecting them with mixed moments)

$$\tau(X_1 X_2 \dots X_n) = \sum_{v \in NC(n)} R_v(X_1, X_2, \dots, X_n), \quad (1)$$

where

Wiktor Ejsmont

Department of Mathematics and Cybernetics, Wrocław University of Economics
wiktor.ejsmont@gmail.com

$$R_\nu(X_1, X_2, \dots, X_n) := \prod_{B \in \nu} R_{|B|}(X_i : i \in B) \quad (2)$$

and $NC(n)$ is the set of all non-crossing partitions of $\{1, 2, \dots, n\}$ (see: [Nica, Speicher 2006; Speicher 1997]). Sometimes we will write $R_k(X) = R_k(X, \dots, X)$.

Definition 1. X is a free normal (Wigner's semicircle law) distribution if $R_k(X) = 0$ for $k > 2$. The Wigner semicircle distribution, is named after the physicist Eugene Wigner. The standardized, i.e. with mean zero and variance one, has density

$$\frac{\sqrt{4-x^2}}{2\pi} 1_{[-2,2]}(x).$$

If $B \subset A$ is a von Neumann subalgebra and A has a trace τ , then there exists a unique conditional expectation from A to B with respect to τ , which we denote by $\tau(\cdot | B)$. This map is weakly continuous, completely positive, identity preserving, contraction and it is characterized by the property that, for any $X \in A$, $\tau(XY) = \tau(\tau(X | B)Y)$ for any $Y \in B$ (see: [Biane 1998; Takesaki 1972]). For fixed $X \in A$ by $\tau(\cdot | X)$ we denote the conditional expectation corresponding to the von Neumann algebra B generated by X and I . The following lemma has been proven in [Bożejko, Bryc 2006].

Lemma 1. Let W be a (self-adjoint) element of the von Neumann algebra A , generated by a self-adjoint $V \in A$. If for all $n \geq 1$ we have $\tau(UV^n) = \tau(WV^n)$, then

$$\tau(U | V) = W. \quad (3)$$

We introduce the notation

- $NC(n)$ is the set of all non-crossing partitions of $\{1, 2, \dots, n\}$,
- $NC^k(m)$ is the set of all non-crossing partitions of $\{1, 2, \dots, m\}$ (where $m \geq k \geq 1$) which have first k elements in the same block (see more: [Ejsmont 2014]).

Let Z be the self-adjoint element of the von Neumann algebra A from the above lemma. We define $c_n^k = c_n^k(Z) = \sum_{\nu \in NC^k(n+k)} R_\nu(Z)$ and the following functions (power series):

$$C^k(z) = \sum_{n=0}^{\infty} c_n^k z^{k+n}, \text{ where } k \geq 1 \quad (4)$$

for sufficiently small $|z| < \varepsilon$ and $z \in C$. The following lemma can be found in [Ejsmont 2014].

Lemma 2. *Let Z be a (self-adjoint) element of the von Neumann algebra A , then*

$$C^{(k)}(z) = M(z)C^{(k+1)}(z) + R_k(Z)z^k M(z), \quad (5)$$

where $k \geq 1$.

Example 1. For $k = 1$, we get:

$$C^{(1)}(z) = M(z) - 1 = M(z)C^{(2)}(z) + R_1(Z)zM(z). \quad (6)$$

In particular, we have the coefficients of the power series $1/M(z)$ (Maclaurin series):

$$\frac{1}{M(z)} = 1 - C^{(2)}(z) - R_1(Z)z \quad (7)$$

for sufficiently small $|z|$.

Similarly, by putting $k = 2$, we obtain:

$$C^{(2)}(z) = M(z)C^{(3)}(z) + R_2(Z)z^2 M(z). \quad (8)$$

Finally, we introduce moment generating function M_X of a random variable X by

$$M_X(z) = \sum_{n=0}^{\infty} \tau(X^n) z^n. \quad (9)$$

Now we present Lemma 4.1 of Bożejko and Bryc [2006], which will be used in the proof of the main theorem.

Lemma 3. *Suppose that X, Y are free, self-adjoint then X, Y have Wigner's semicircle law if and only if the moment generating function $M(z)$ for $X + Y$ satisfies the following quadratic equation*

$$2z^2 M^2(z) - M(z) + 1 = 0. \quad (10)$$

2. Characterization theorem

The main result of this paper is the following characterization of Wigner's semicircle law.

Theorem 1. *Suppose that X, Y are free, self-adjoint, non-degenerate, centered ($\tau(X) = \tau(Y) = 0$) and $\tau(X^2) = \tau(Y^2) = 1$ which have the same distribution. Additionally, we assume that $R_3(X) = R_3(Y) = 0$. Then X and Y have Wigner's semicircle law if and only if*

$$\tau\left((X - Y)(X + Y)^2(X - Y) \mid (X + Y)\right) = 4I. \quad (11)$$

Proof. \Leftarrow : Let us suppose now that the equality (11) holds. Multiplying (11) by $(Y + Y)^n$ for $n \geq 0$ and applying $\tau(\cdot)$ we obtain

$$\begin{aligned} \tau((X - Y)(X + Y)^2(X - Y)(X + Y)^n) = \\ \sum_{v \in NC(n+4)} R_v(X - Y, X + Y, X + Y, X - Y, \underbrace{X + Y, X + Y, \dots, X + Y}_{n\text{-times}}) = \\ 4\tau((X + Y)^n). \end{aligned}$$

This follows from the following consideration. Let us look more closely at the second sum from the last equation. We have that either the first and the fourth elements are in different blocks, or they are in the same block. In the first case, the second sum (from the last equation) vanishes because we have

$$R_k(X - Y, X + Y, X + Y, \dots, X + Y) = R_k(X) - R_k(Y) = 0. \quad (12)$$

On the other hand, if they are in the same block, the sum disappears if the first or third element are in the same block separately because we have that $\tau(X + Y) = 0$. So, we have

$$\begin{aligned} \tau\left((X - Y)(X + Y)^2(X - Y)(X + Y)^n\right) = \\ \sum_{v \in NC^4(n+4)} R_v(X + Y) + 2 \sum_{v \in NC^2(n+2)} R_v(X + Y) = 4\tau((X + Y)^n). \end{aligned} \quad (13)$$

This equation is equivalent to

$$4z^4 M(z) = C^4(z) + 2z^2 C^2(z). \quad (14)$$

Using Lemma 2 for $k = 1, 2, 3$ we obtain equation

$$R_3(X + Y)z^3M^3(z) = (M(z) - 1 - 2z^2M^2(z))(1 + 2z^2M^2(z)). \quad (15)$$

Thus, if $R_3(X + Y) = 0$, then we have found two solutions

$$1 + 2z^2M^2(z) = 0 \quad (16)$$

or

$$M(z) - 1 - 2z^2M^2(z) = 0 \quad (17)$$

but the first solution does not correspond to the probability measure. Part $M(z) - 1 - 2z^2M^2(z)$ corresponds to the moment generating function of Wigner's semicircle law, which by **Lemma 3** implies statement. \Rightarrow : Suppose that X and Y have Wigner's semicircle law. Then we have

$$\sum_{v \in NC^4(n+4)} R_v(X + Y) = 0.$$

So, from $R_2(X + Y) = 2$ we see

$$\begin{aligned} \tau((X - Y)(X + Y)^2(X - Y)(X + Y)^n) = \\ 2 \sum_{v \in NC^2(n+2)} R_v(X + Y) = 4 \sum_{v \in NC(n)} R_v(X + Y) = 4\tau((X + Y)^n). \end{aligned} \quad (18)$$

Now **Lemma 1** implies equation (11).

References

- Biane P. (1998). *Processes with free increments*. *Mathematische Zeitschrift* 227(1). Pp. 143-174.
- Bożejko M., Bryc W. (2006). *On a class of free Lévy laws related to a regression problem*. *Journal of Functional Analysis* 236. Pp. 59-77.
- Ejsmont W. (2014). *Characterizations of some free random variables by properties of conditional moments of third degree polynomials*. *Journal of Theoretical Probability* 27 (3). Pp. 915-931.
- Nica A., Speicher R. (2006). *Lectures on the Combinatorics of Free Probability*. London Mathematical Society Lecture Notes Series 365. Cambridge University Press.

- Speicher R. (1997). *Free probability theory and non-crossing partitions*. Séminaire Lotharingien de Combinatoire B39c. Pp. 38. (electronic)
<http://www.emis.ams.org/journals/SLC/wpapers/s39speicher.pdf>.
- Takesaki M. (1972). *Conditional expectations in von Neumann algebras*. Journal of Functional Analysis 9. Pp. 306-321.