

**ON SOME INTEGRAL INEQUALITIES
FOR (h, m) -CONVEX FUNCTIONS****Marian Matloka**

Abstract. In this paper we establish several Hadamard type inequalities for (h, m) convex functions.

Keywords: Hadamard's inequality, convex function.

JEL Classification: C02.

DOI: 10.15611/me.2013.9.05.

1. Introduction

A function $f: I \rightarrow R$, $I \subseteq R$ is an interval, said to be a convex function on I if

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y) \quad (1.1)$$

holds for all $x, y \in I$ and $t \in [0, 1]$. If the reversed inequality in (1.1) holds, then f is concave.

Many important inequalities have been established for the class of convex functions, but the most famous is Hermite-Hadamard's inequality. This double inequality is stated as:

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2} \quad (1.2)$$

where $f: [a, b] \rightarrow R$, is a convex function. The above inequalities are in reversed order if f is a concave function.

Marian Matloka

Department of Applied Mathematics, Poznań University of Economics, Al. Niepodległości 10,
61-875 Poznań, Poland.

E-mail: marian.matloka@ue.poznan.pl

In 1978, Breckner introduced the s -convex function as a generalization of the convex function (Breckner 1978). Such a function is defined in the following way: a function $f : [0, \infty] \rightarrow R$ is said to be s -convex in the second sense if

$$f(tx + (1-t)y) \leq t^s f(x) + (1-t)^s f(y) \quad (1.3)$$

holds for all $x, y \in [0, \infty]$, $t \in [0, 1]$ and for fixed $s \in [0, 1]$.

In (Dragomir, Fitzpatrick 1999) Dragomir and Fitzpatrick proved the following variant of Hermite-Hadamard's inequality which holds for s -convex functions in the second sense.

$$2^{s-1} f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{s+1}. \quad (1.4)$$

In the paper (Varošanec 2007) a large class of non-negative functions, the so-called h -convex functions, is considered. This class contains several well-known classes of functions such as non-negative convex functions and s -convex in the second sense. This class is defined in the following way: a non-negative function $f : I \rightarrow R$, $I \subseteq R$ is an interval, called h -convex if

$$f(tx + (1-t)y) \leq h(t)f(x) + h(1-t)f(y) \quad (1.5)$$

holds for all $x, y \in I$, $t \in (0, 1)$, where $h : J \rightarrow R$ is a non-negative function, $h \not\equiv 0$ and J is an interval, $(0, 1) \subseteq J$.

In (Sarıkaya, Saglam, Yildirim 2008) the authors proved that for h -convex function the following variant of Hadamard inequality is fulfilled

$$\frac{1}{2h\left(\frac{1}{2}\right)} f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq [f(a) + f(b)] \cdot \int_0^1 h(t) dt \quad (1.6)$$

In 1988, Weir and Mond (1998) introduced the preinvex function. Such a function is defined in the following way: a function f on the invex set X is said to be preinvex with respect to η , if

$$f(u + t\eta(v, u)) \leq (1-t)f(u) + tf(v) \quad (1.7)$$

for each $u, v \in X$ and $t \in [0, 1]$, where $\eta : X \times X \rightarrow R$.

Noor in (Noor 2009) proved the Hermite-Hadamard inequality for the preinvex functions:

$$f\left(a + \frac{1}{2}\eta(b,a)\right) \leq \frac{1}{\eta(b,a)} \int_a^{a+\eta(b,a)} f(x) dx \leq \frac{f(a) + f(b)}{2}. \quad (1.8)$$

Matłoka introduced in (Matłoka 2013) the h -preinvex function in the following way: The non-negative function f on the invex set X is said to be h -preinvex with respect to η , if

$$f(u + t\eta(v,u)) \leq h(1-t)f(u) + h(t)f(v) \quad (1.9)$$

for each $u, v \in X$ and $t \in [0,1]$.

In the same paper Matłoka proved the Hermite-Hadamard inequality for the h -preinvex functions:

$$\begin{aligned} \frac{1}{2h\left(\frac{1}{2}\right)} f\left(a + \frac{1}{2}\eta(b,a)\right) &\leq \frac{1}{\eta(b,a)} \int_a^{a+\eta(b,a)} f(x) dx \\ &\leq [f(a) + f(b)] \int_0^1 h(t) dt. \end{aligned} \quad (1.10)$$

Toader (1985) defined m -convexity in the following way: the function $f: [0,b] \rightarrow R$, $b > 0$, is said to be m -convex, where $m \in [0,1]$, if

$$f(tx + m(1-t)y) \leq tf(x) + m(1-t)f(y) \quad (1.11)$$

for all $x, y \in [0,b]$ and $t \in [0,1]$.

In (Dragomir, Toader 1993) the authors proved the following Hadamard type inequality for m -convex functions:

$$\frac{1}{b-a} \int_a^b f(x) dx \leq \min \left\{ \frac{f(a) + m f\left(\frac{b}{m}\right)}{2}, \frac{f(b) + m f\left(\frac{a}{m}\right)}{2} \right\}. \quad (1.12)$$

In this paper we introduce the concept of the (h, m) -convex function. The main purpose of this paper is to establish new inequalities of the class of (h, m) -convex functions.

2. Inequalities for (h, m) -convex functions

Definition 2.1. Let $h: [0,1] \rightarrow R$ be a nonnegative function, $h \neq 0$. The non-negative function $f: [0,b] \rightarrow R$, $b > 0$, is said to be (h, m) -convex, where $m \in [0,1]$, if we have

$$f(tx + m(1-t)y) \leq h(t)f(x) + mh(1-t)f(y)$$

for all $x, y \in [0,b]$ and $t \in [0,1]$.

If the above inequality is reversed, then f is said to be (h, m) -preconcave.

Note that if $h(t) = t$ then the f above definition reduces to the definition of m -convex function.

Definition 2.2. The function $f: [0,b] \rightarrow R$, $b > 0$, is said to be (h, m) -logarithmic convex, where $m \in [0,1]$, if

$$\log f(tx + m(1-t)y) \leq h(t)\log f(x) + mh(1-t)\log f(y)$$

for all $x, y \in [0,b]$, $t \in [0,1]$, where $f(\cdot) > 0$.

If the above inequality is reversed, then f is said to be (h, m) -logarithmic concave.

From now on we suppose that all the integrals of function h considered below exist.

Theorem 2.1. Let $f: [0,\infty] \rightarrow R$ be a (h, m) -convex function with $m \in (0,1]$. If $0 \leq a < mb < \infty$ and $f \in L^1([a,mb])$, $h \in L^1([0,1])$ then

$$\frac{1}{mb-a} \int_a^{mb} f(x) dx \leq [f(a) + mf(b)] \cdot \int_0^1 h(t) dt. \quad (2.1)$$

Proof. From the (h, m) -convexity of f we have

$$f(ta + m(1-t)b) \leq h(t)f(a) + mh(1-t)f(b).$$

Thus by integrating over $[0,1]$ we obtain

$$\int_0^1 f(ta + m(1-t)b) dt \leq f(a) \int_0^1 h(t) dt + mf(b) \int_0^1 h(1-t) dt.$$

Since,

$$\int_0^1 f(ta + m(1-t)b) dt = \frac{1}{mb-a} \int_a^{mb} f(x) dx$$

then

$$\frac{1}{mb-a} \int_a^{mb} f(x) dx \leq [f(a) + mf(b)] \cdot \int_0^1 h(t) dt$$

which completes the proof.

Remark 2.1.

– if $m = 1$ and $h(t) = t$ then inequality (2.1) reduces to the right hand of the Hermite-Hadamard inequality for convex function.

– if $m = 1$ and $h(t) = t^s$, $s \in [0, 1]$ then we obtain the right hand of a variant of the Hadamard inequality (1.4) for s -convex function in the second sense.

– if $m = 1$ then inequality (2.1) reduces to the right hand of the Hadamard inequality (1.6) for h -convex function (see Sarikaya et. al. 2008).

In an analogous way we can prove the following inequality for (h, m) -logarithmic convex function

$$\frac{1}{mb-a} \int_a^{mb} \log f(x) dx \leq [\log f(a) + m \log f(b)] \cdot \int_0^1 h(t) dt. \quad (2.2)$$

Theorem 2.2. Let f be a (h_1, m) -convex and g a (h_2, m) -convex functions such that $f \cdot g \in L^1([a, b])$ and $h_1 \cdot h_2 \in L^1([0, 1])$. Then the following inequality holds:

$$\begin{aligned} \frac{1}{mb-a} \int_a^{mb} f(x)g(x) dx &\leq [f(a)g(a) + m^2 f(b)g(b)] \cdot \int_0^1 h_1(t)h_2(t) dt \\ &+ m[f(a)g(b) + f(b)g(a)] \cdot \int_0^1 h_1(t)h_2(1-t) dt. \end{aligned} \quad (2.3)$$

Proof. Using the fact that f and g are (h_1, m) -convex and (h_2, m) -convex respectively we have

$$\begin{aligned}
& (f \cdot g)(ta + m(1-t)b) \\
& \leq [h_1(t)f(a) + mh_1(1-t)f(b)] \cdot [h_2(t)g(a) + mh_2(1-t)g(b)] \\
& = h_1(t)h_2(t)f(a)g(a) + m^2h_1(1-t)h_2(1-t)f(b)g(b) \\
& \quad + mh_1(t)h_2(1-t)f(a)g(b) + mh_1(1-t)h_2(t)f(b)g(a).
\end{aligned}$$

Thus, by integrating with respect to t over $[0, 1]$, we obtain

$$\begin{aligned}
\int_0^1 (f \cdot g)(ta + m(1-t)b) dt & \leq [f(a)g(a) + m^2f(b)g(b)] \int_0^1 h_1(t)h_2(t) dt \\
& \quad + m[f(a)g(b) + f(b)g(a)] \int_0^1 h_1(t)h_2(1-t) dt.
\end{aligned}$$

Since

$$\int_0^1 (f \cdot g)(ta + m(1-t)b) dt = \frac{1}{mb-a} \int_a^{mb} f(x)g(x) dx$$

then we obtain the inequality (2.3).

Theorem 2.3. Let f be a (h_1, m_1) -convex and g a (h_2, m_2) -convex functions such that $f \cdot g \in L^1([a, mb])$ and $h_1 \cdot h_2 \in L^1([0, 1])$. Then the following inequality holds:

$$\begin{aligned}
& \frac{1}{mb-a} \int_a^{mb} f(x)g(x) dx \\
& \leq \min \left\{ M_1 \cdot \int_0^1 h_1(t)h_2(t) dt + M_2 \int_0^1 h_1(t)h_2(1-t) dt M_3 \right. \\
& \quad \left. \cdot \int_0^1 h_1(t)h_2(t) dt + M_4 \int_0^1 h_1(t)h_2(1-t) dt \right\}, \tag{2.4}
\end{aligned}$$

where

$$M_1 = f(a)g(a) + m_1m_2 f\left(\frac{b}{m_1}\right)g\left(\frac{b}{m_2}\right),$$

$$M_2 = m_2 f(a) g\left(\frac{b}{m_2}\right) + m_1 f\left(\frac{b}{m_1}\right) g(a),$$

$$M_3 = m_1 m_2 f\left(\frac{a}{m_1}\right) g\left(\frac{a}{m_2}\right) + f(b) g(b),$$

$$M_4 = m_1 f\left(\frac{a}{m_1}\right) g(b) + m_2 f(b) g\left(\frac{a}{m_2}\right).$$

Proof. Using the fact that f i g are (h_1, m_1) -convex and (h_2, m_2) -convex respectively we have

$$\begin{aligned} f\left(ta + (1-t)b\right) \cdot g\left(ta + (1-t)b\right) &= f\left(ta + m_1(1-t)\frac{b}{m_1}\right) \\ \cdot g\left(ta + m_2(1-t)\frac{b}{m_2}\right) &\leq \left[h_1(t) f(a) + m_1 h_1(1-t) f\left(\frac{b}{m_1}\right) \right] \\ &\quad \cdot \left[h_2(t) g(a) + m_2 h_2(1-t) g\left(\frac{b}{m_2}\right) \right] \\ &= h_1(t) f(a) h_2(t) g(a) + m_1 m_2 h_1(1-t) h_2(1-t) f\left(\frac{b}{m_1}\right) g\left(\frac{b}{m_2}\right) \\ &\quad + m_2 h_1(t) f(a) h_2(1-t) g\left(\frac{b}{m_2}\right) + m_1 h_1(1-t) f\left(\frac{b}{m_1}\right) h_2(t) g(a). \end{aligned}$$

Integrating both sides of the above inequality over $[0,1]$ we obtain

$$\begin{aligned} \int_0^1 f\left(ta + (1-t)b\right) g\left(ta + (1-t)b\right) dt &= \frac{1}{b-a} \int_a^b f(x) g(x) dx \\ &\leq \left[f(a) g(a) + m_1 m_2 f\left(\frac{b}{m_1}\right) g\left(\frac{b}{m_2}\right) \right] \int_0^1 h_1(t) h_2(t) dt \\ &\quad + \left[m_2 f(a) g\left(\frac{b}{m_2}\right) + m_1 f\left(\frac{b}{m_1}\right) g(a) \right] \int_0^1 h_1(t) h_2(1-t) dt. \end{aligned}$$

Analogously we obtain

$$\begin{aligned} \frac{1}{b-a} \int_a^b f(x)g(x) \leq & \left[m_1 m_2 f\left(\frac{a}{m_1}\right) g\left(\frac{a}{m_2}\right) + f(b)g(b) \right] \int_0^1 h_1(t)h_2(t)dt \\ & + \left[m_1 f\left(\frac{a}{m_1}\right) g(b) + m_2 f(b)g\left(\frac{a}{m_2}\right) \right] \int_0^1 h_1(t)h_2(1-t)dt \end{aligned}$$

which completes the proof.

Let us note that from the inequality (2.2) it follows the following inequality for (h_1, m) -log-convex function f and (h_2, m) -log-convex function g :

$$\begin{aligned} & \frac{1}{mb-a} \int_a^{mb} \log(f(x) \cdot g(x)) dx \\ & \leq [\log f(a) + m \log f(b)] \int_0^1 h_1(t) dt \\ & \quad + [\log g(a) + m \log g(b)] \int_0^1 h_2(t) dt. \end{aligned}$$

Moreover, if f is (h_1, m) -log-convex and g is (h_2, m) -log-concave then from the some inequality it follows that

$$\begin{aligned} \frac{1}{mb-a} \int_a^{mb} \log \frac{f(x)}{g(x)} dx \leq & [\log f(a) + m \log f(b)] \cdot \int_0^1 h_1(t) dt \\ & - [\log g(a) + m \log g(b)] \cdot \int_0^1 h_2(t) dt. \end{aligned}$$

Using the technique and ideas of Bakula, Özdemir and Pečarić (2008, Theorem 2.1), one can prove the following theorem.

Theorem 2.4. Let I be an open real interval such that $[0, \infty) \subset I$. Let $f : I \rightarrow \mathbb{R}$ be a differentiable function on I such that $f' \in L^1([a, b])$, where $0 \leq a < b < \infty$. If $|f'|^q$ is (h, m) -convex on $[a, b]$ for some fixed $m \in (0, 1]$ and $q \in [1, \infty]$, then

$$\begin{aligned} & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \frac{b-a}{2} \cdot \left[\int_{\frac{1}{2}}^1 h(1-t)(2t-1) dt + \int_{\frac{1}{2}}^1 h(t)(2t-1) dt \right]^{\frac{1}{q}} \\ & \cdot \min \left\{ \left(|f'(a)|^q + m \left| f' \left(\frac{b}{m} \right) \right|^q \right)^{\frac{1}{q}}, \left(m \left| f' \left(\frac{a}{m} \right) \right|^q + \left| f' \left(\frac{b}{m} \right) \right|^q \right)^{\frac{1}{q}} \right\}. \end{aligned}$$

Proof. First let us note that for a differentiable mapping f such that $f' \in L^1([a, b])$ the following equation holds

$$\frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx = \frac{b-a}{2} \int_0^1 (1-2t) f'(ta + (1-t)b) dt.$$

First let us suppose that $q = 1$. Then from the above equation we have

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{b-a}{2} \int_0^1 |1-2t| \cdot |f'(ta + (1-t)b)| dt.$$

Since $|f'|$ is (h, m) -convex on $[a, b]$ we know that

$$\left| f'(ta + (1-t)b) \right| = \left| f' \left(ta + m(1-t) \frac{b}{m} \right) \right| \leq h(t) |f'(a)| + mh(1-t) \left| f' \left(\frac{b}{m} \right) \right|,$$

hence

$$\begin{aligned} & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \frac{b-a}{2} \int_0^1 |1-2t| \cdot \left[h(t) |f'(a)| + mh(1-t) \left| f' \left(\frac{b}{m} \right) \right| \right] dt = \end{aligned}$$

$$\begin{aligned}
&= \frac{b-a}{2} \left\{ \int_0^{\frac{1}{2}} (1-2t) \left[h(t) |f'(a)| + mh(1-t) \cdot \left| f' \left(\frac{b}{m} \right) \right| \right] dt \right. \\
&\quad \left. + \int_{\frac{1}{2}}^1 (2t-1) \left[h(t) |f'(a)| + mh(1-t) \left| f' \left(\frac{b}{m} \right) \right| \right] dt \right\} \\
&= \frac{b-a}{2} \left[|f'(a)| + m \left| f' \left(\frac{b}{m} \right) \right| \right] \cdot \left[\int_{\frac{1}{2}}^1 h(1-t)(2t-1) dt + \int_{\frac{1}{2}}^1 h(t)(2t-1) dt \right],
\end{aligned}$$

where we have used the fact that

$$\int_0^{\frac{1}{2}} (1-2t)h(1-t) dt = \int_{\frac{1}{2}}^1 h(t)(2t-1) dt$$

and

$$\int_0^{\frac{1}{2}} (1-2t)h(t) dt = \int_{\frac{1}{2}}^1 h(1-t)(2t-1) dt.$$

Analogously we obtain

$$\begin{aligned}
&\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\
&= \frac{b-a}{2} \cdot \left[m \left| f' \left(\frac{a}{m} \right) \right| + |f'(b)| \right] \cdot \left[\int_{\frac{1}{2}}^1 h(1-t)(2t-1) dt + \int_{\frac{1}{2}}^1 h(t)(2t-1) dt \right],
\end{aligned}$$

which completes the proof for $q = 1$.

Suppose now that $q > 1$. Since $|f'|^q$ is (h, m) -convex on $[a, b]$

$$\left| f'(ta + (1-t)b) \right|^q \leq h(t) |f'(b)|^q + mh(1-t) \cdot \left| f' \left(\frac{b}{m} \right) \right|^q$$

hence using the well-known Hölder inequality we obtain

$$\begin{aligned} & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ &= \frac{b-a}{2} \left(\int_0^1 |1-2t| dt \right)^{\frac{q-1}{q}} \cdot \left(\int_0^1 |1-2t| \cdot \left| f' \left(ta + m(1-t) \frac{b}{m} \right) \right|^q dt \right)^{\frac{1}{q}} \\ &\leq \frac{b-a}{2} \cdot \left(\int_{\frac{1}{2}}^1 h(1-t)(2t-1) dt + \int_{\frac{1}{2}}^1 h(t)(2t-1) dt \right) \cdot \left[|f'(a)|^q + m \left| f' \left(\frac{b}{m} \right) \right|^q \right]^{\frac{1}{q}} \end{aligned}$$

and analogously

$$\begin{aligned} & \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ &\leq \frac{b-a}{2} \cdot \left(\int_{\frac{1}{2}}^1 h(1-t)(2t-1) dt + \int_{\frac{1}{2}}^1 h(t)(2t-1) dt \right)^{\frac{1}{q}} \cdot \left[m \left| f' \left(\frac{a}{m} \right) \right|^q + |f'(b)|^q \right]^{\frac{1}{q}} \end{aligned}$$

which completes the proof.

Using the identity

$$f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(x) dx = \frac{1}{b-a} \int_a^b S(x) f'(x) dx,$$

where

$$S(x) = \begin{cases} x-a, & x \in \left[a, \frac{a+b}{2} \right] \\ x-b, & x \in \left[\frac{a+b}{2}, b \right] \end{cases}$$

(see (Pearce, Pečarić 2000, Theorem 2)) we can prove the following theorem.

Theorem 2.5. Let I be an open real interval such that $[0, \infty) \subset I$. Let $f : I \rightarrow \mathbb{R}$ be a differentiable function on I such that $f' \in L^1([a, b])$, where $0 \leq a < b < \infty$. If $|f'|$ is (h, m) -convex on $[a, b]$ for some fixed $m \in [0, 1]$, then

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq (b-a) \left(\int_0^{\frac{1}{2}} th(t) dt + \int_0^{\frac{1}{2}} th(1-t) dt \right) \min \left\{ |f'(a)| + m \left| f'\left(\frac{b}{m}\right) \right|; m \left| f'\left(\frac{a}{m}\right) \right| + |f'(b)| \right\}. \end{aligned}$$

Proof.

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \frac{1}{b-a} \left[\int_a^{\frac{a+b}{2}} (x-a) |f'(x)| dx + \int_{\frac{a+b}{2}}^b (b-x) |f'(x)| dx \right] \\ & = (b-a) \left[\int_0^{\frac{1}{2}} t |f'(ta + (1-t)b)| dt + \int_{\frac{1}{2}}^1 (1-t) |f'(ta + (1-t)b)| dt \right] \\ & \leq (b-a) \left[\int_0^{\frac{1}{2}} t \left(h(t) |f'(a)| + mh(1-t) \left| f'\left(\frac{b}{m}\right) \right| \right) dt \right. \\ & \quad \left. + \int_{\frac{1}{2}}^1 (1-t) \left(h(t) |f'(a)| + mh(1-t) \left| f'\left(\frac{b}{m}\right) \right| \right) dt \right] \\ & = (b-a) \left(|f'(a)| + m \left| f'\left(\frac{b}{m}\right) \right| \right) \cdot \left(\int_0^{\frac{1}{2}} th(t) dt + \int_0^{\frac{1}{2}} th(1-t) dt \right) \end{aligned}$$

and analogously

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq (b-a) \left(m \left| f'\left(\frac{a}{m}\right) \right| + m |f'(b)| \right) \cdot \left(\int_0^{\frac{1}{2}} th(t) dt + \int_0^{\frac{1}{2}} th(1-t) dt \right) \end{aligned}$$

which completes the proof.

Now, let us note that it can be easy to prove the following two lemmas.

Lemma 2.1. Let $f : I \rightarrow R$, $I \subset R$, be a differentiable mapping on I° , and $a, b \in I$, $m \in [0, 1]$ and $a < mb$. If $f' \in L^1([a, mb])$, then

$$\frac{f(a) + f(mb)}{2} - \frac{1}{mb-a} \int_a^{mb} f(x) dx = \frac{mb-a}{2} \int_0^1 (1-2t) f'(ta + m(1-t)b) dt.$$

Lemma 2.2. Let $f : I \rightarrow R$, $I \subset R$, be a differentiable mapping on I° , and $a, b \in I$, $m \in [0, 1]$ and $a < mb$. If $f' \in L^1([a, mb])$, then

$$\begin{aligned} & \frac{1}{mb-a} \int_a^{mb} f(x) dx - f\left(\frac{a+mb}{2}\right) \\ & = (mb-a) \left[\int_0^{\frac{1}{2}} t f'(ta + m(1-t)b) dt + \int_{\frac{1}{2}}^1 (t-1) f'(ta + m(1-t)b) dt \right]. \end{aligned}$$

Theorem 2.6. Let $f : I \rightarrow R$, be a differentiable function on I° that $f' \in L^1([a, mb])$, where $a, b \in I$, $m \in [0, 1]$ and $a < mb$. If $|f'|$ is (h, m) -convex function, then we have

$$\begin{aligned} & \left| \frac{f(a) + f(mb)}{2} - \frac{1}{mb-a} \int_a^{mb} f(x) dx \right| \\ & \leq \frac{mb-a}{2} \left[|f'(a)| + m |f'(b)| \right] \cdot \left[\int_{\frac{1}{2}}^1 h(1-t)(2t-1) dt + \int_{\frac{1}{2}}^1 h(t)(2t-1) dt \right]. \end{aligned}$$

Proof. Using Lemma 2.1 and the (h, m) -convexity of $|f'|$ we have

$$\begin{aligned}
& \left| \frac{f(a) + f(mb)}{2} - \frac{1}{mb-a} \int_a^{mb} f(x) dx \right| \\
& \leq \frac{mb-a}{2} \int_0^1 |1-2t| (h(t)|f'(a)| + mh(1-t)|f'(b)|) dt \\
& = \frac{mb-a}{2} \left[\int_0^{\frac{1}{2}} (|1-2t|h(t)|f'(a)| + mh(1-t)(|f'(b)|)) dt \right. \\
& \quad \left. + \int_{\frac{1}{2}}^1 (2t-1)(h(t)|f'(a)| + mh(1-t)|f'(b)|) dt \right] \\
& = \frac{mb-a}{2} \left[|f'(a)| \int_0^{\frac{1}{2}} (1-2t)h(t) dt + m|f'(b)| \int_0^{\frac{1}{2}} (1-2t)h(1-t) dt \right. \\
& \quad \left. + |f'(a)| \int_{\frac{1}{2}}^1 (2t-1)h(t) dt + m|f'(b)| \int_{\frac{1}{2}}^1 (2t-1)h(1-t) dt \right] \\
& = \frac{mb-a}{2} \left[|f'(a)| \int_{\frac{1}{2}}^1 h(1-t)(2t-1) dt + m|f'(b)| \int_{\frac{1}{2}}^1 h(t)(2t-1) dt \right. \\
& \quad \left. + |f'(a)| \int_{\frac{1}{2}}^1 h(t)(2t-1) dt + m|f'(b)| \int_{\frac{1}{2}}^1 h(1-t)(2t-1) dt \right] \\
& = \frac{mb-a}{2} [|f'(a)| + m|f'(b)|] \left[\int_{\frac{1}{2}}^1 h(1-t)(2t-1) dt + \int_{\frac{1}{2}}^1 h(t)(2t-1) dt \right],
\end{aligned}$$

which completes the proof.

Theorem 2.7. Let $f : I \rightarrow R$, be a differentiable function on I° , with $a, b \in I$, $m \in [0, 1]$ and $a < mb$. If $|f'|$ is (h, m) -convex, then we have

$$\begin{aligned} & \left| \frac{1}{mb-a} \int_a^{mb} f(x) dx - f\left(\frac{a+mb}{2}\right) \right| \\ &= (mb-a) \left[|f'(a)| + m|f'(b)| \right] \cdot \left[\int_0^{\frac{1}{2}} th(t) dt + \int_0^{\frac{1}{2}} th(1-t) dt \right]. \end{aligned}$$

Proof. Using Lemma 2.2 and the (h, m) -convexity of $|f'|$, it follows that

$$\begin{aligned} & \left| \frac{1}{mb-a} \int_a^{mb} f(x) dx - f\left(\frac{a+mb}{2}\right) \right| \\ & \leq (mb-a) \left[\int_0^{\frac{1}{2}} |t| |f'(ta+m(1-t)b)| dt + \int_{\frac{1}{2}}^1 |t-1| |f'(ta+m(1-t)b)| dt \right] \\ & \leq (mb-a) \left[\int_0^{\frac{1}{2}} |t| (|h(t)| |f'(a)| + mh(1-t) |f'(b)|) dt \right. \\ & \quad \left. + \int_{\frac{1}{2}}^1 |t-1| (|h(t)| |f'(a)| + mh(1-t) |f'(b)|) dt \right] \\ & = (mb-a) \left[|f'(a)| \int_0^{\frac{1}{2}} th(t) dt + m|f'(b)| \int_0^{\frac{1}{2}} th(1-t) dt \right. \\ & \quad \left. + |f'(a)| \int_0^{\frac{1}{2}} th(1-t) dt + m|f'(b)| \int_0^{\frac{1}{2}} th(t) dt \right] \\ & = (mb-a) \left[|f'(a)| + m|f'(b)| \right] \cdot \left[\int_0^{\frac{1}{2}} th(t) dt + \int_0^{\frac{1}{2}} th(1-t) dt \right] \end{aligned}$$

which completes the proof.

References

- Bakula M.K., Özdemir M.E., Pečarić J. (2008). *Hadamard type inequalities for m -convex and (α, m) -convex functions*. Journal of Inequalities in Pure and Applied Mathematics 9.
- Breckner W.W. (1978). *Stetigkeitsanssagen für eine Klasse verallgemeinerter Konvexer Funktionen in topologischen linearen Räumen*, Publications de l'Institut Mathématique. 23 (37). Pp. 13-20.
- Dragomir S.S., Fitzpatrick S. (1999). *The Hadamard's inequality for s – convex functions in the second sense*. Demonstration Mathematics 32 (4). Pp. 687-696.
- Dragomir S.S., Toader G. (1993). *Some inequalities for m – convex functions*. Studia Universitatis Babeş-Bolyai Mathematica 38. Pp. 21-28.
- Kirmaci U. S., Bakula M. K., Özdemir M. E., Pečarić J. (2007). *Hadamard-type inequalities for s -convex functions*. Applied Mathematics and Computation 193. Pp. 26-35.
- Matloka M. (2013). *On some Hadamard-type inequalities for (h_1, h_2) -preinvex functions on the co-ordinates*. Journal of Inequalities and Applications. doi: 10.1186/1029-242X-2013-227.
- Noor M.S. (2009). *Hadamard integral inequalities for product of two preinvex functions*. Nonlinear Analysis Forum 14. Pp. 167-173.
- Pearce C.E.M., Pečarić J. (2000). *Inequalities for differentiable mappings with application to special means and quadrature formulae*. Applied Mathematical Letter 13. Pp. 51-55.
- Sarikaya M.Z., Saglam A., Yildirim H. (2008). *On some Hadamard-type inequalities for h -convex functions*. Journal of Mathematical Inequalities 2. Pp. 335-341.
- Toader G. (1985). *Some generalizations of the convexity*. Proceedings of the Colloquium on Approximation and Optimization. University Cluj-Napoca. Pp. 329-338.
- Weir T., Mond B. (1988). *Preinvex functions in multiobjective optimization*. Journal of Mathematical Analysis and Application 136. Pp. 29-38.
- Varošanec S. (2007). *On h -convexity*. Journal of Mathematical Analysis and Application 326. Pp. 303-311.